Chpt the 3rd

Exponential and Logarithmic Functions

3.4 Exponential and Logarithmic Equations

128

Chpt 3-4

Homework

3-4 p432 11, 13, 19, 31, 35, 41, 43, 53, 55, 63, 69, 77, 81, 85, 89



1701

Chpt 3-4



Use like bases to solve exponential equations.
 Use logarithms to solve exponential equations.
 Use the definition of a logarithm to solve logarithmic equations.
 Use the one-to-one property of logarithms to solve logarithmic equations.
 Solve applied problems involving exponential and logarithmic equations.

xponential and logarithmic equations. 3/28

C-1701

Solving Expnential Equations

There are two two primary methods for solving exponential equations:

- I. Isolate the exponential expression.
- 2. Use the one-to-one property.

There are also two primary ways of solving logarithmic equations.

- reformable and the logarithmic expression, then write the equation in equivalent exponential form.
- 2. Get a single logarithmic expression with the same base on each side of the equation; then use the one-to-one property.
- One to one properties

 \ll a×=aY if and only if x = y.

 \ll log_a = log_a if and only if $\chi = \gamma$.



Note: In both cases the base must be the same.



Solving Exponential Equations by Expressing Each Side as a Power of the Same Base


Solving Using Exponential Equations with common base.

 \sim If $b^m = b^n$ then m = n.

1. Rewrite the equation in the form $b^m = b^n$. The base (b) is the same in both expressions.

2. Set m = n.

3. Solve the equation.






Example: Solving Exponential Equations*



- $5^{3x-6} = 125$
- $5^{3x-6} = 5^{3}$
- 3x 6 = 3
 - x = 3

\checkmark The solution set is $\{3\}$





- $8^{x+2} = 4^{x-3}$
- $(2^3)^{x+2} = (2^2)^{x-3}$
- $2^{3(x+2)} = 2^{2(x-3)}$
- 3(x + 2) = 2(x 3)
 - 3x + 6 = 2x 6
 - x = -12
- \sim The solution set is $\{-12\}$



Example: Solving Exponential Equations*



$$e^{x+1} = \frac{1}{e}$$

$$e^{x+1} = e^{-1}$$

x + 1 = -1

\sim The solution set is $\{-2\}$





$$9^{x} = \frac{1}{\sqrt[3]{3}}$$

$$9^{x} = \frac{1}{\frac{1}{3^{\frac{1}{3}}}} \qquad (3^{2})^{x} = 3^{-\frac{1}{3}}$$

$$9^{x} = 3^{-\frac{1}{3}} \qquad 3^{2x} = 3^{-\frac{1}{3}}$$

$$2x = -\frac{1}{3}$$
The solution set is $\left\{-\frac{1}{6}\right\}$

Using Logarithms to Solve Exponential Equations

Solving by taking logarithm of both sides.

- 1. Isolate the exponential expression.
- 2. It the base of the exponent is 10, take the common logarithm of both sides of the equation. Use the natural logarithm on both sides of the equation for bases other than 10.
- 3. Simplify using: $lnb^{\times} = xlnb$ or $lne^{\times} = x$ or $log10^{\times} = x$.
- 4. Solve for the variable.



STUDY TIP

Remember that the natural logarithmic function has a base of *e*.



Solving Exponential Equations



$$x = \frac{\ln 134}{\ln 5} \approx 3.043$$

The solution set is
$$\left\{ \frac{\ln 134}{\ln 5} \right\}$$



The solution set is log8000.

- x ≈ 3.903
- x = log8000
- xlog10 = log8000
- $log10^{\times} = log8000$
- $10^{\times} = 8000$



Solving an Exponential Equation



 $3^{2x-1} = 7^{x+1}$ $ln3^{2x-1} = ln7^{x+1}$ $(2x-1)\ln 3 = (x+1)\ln 7$ 2xln3 - ln3 = xln7 + ln72xln3 - xln7 = ln3 + ln7 $x(2\ln 3 - \ln 7) = \ln 3 + \ln 7$



$$x = \frac{\ln 3 + \ln 7}{2\ln 3 - \ln 7}$$

$$x = \frac{\ln 21}{\ln 9 - \ln 7} = \frac{\ln 21}{\ln \frac{9}{7}} \approx 12.1144$$

The solution set is
$$\left\{\frac{\ln 21}{\ln \frac{9}{7}}\right\}$$

Solving Exponential Equations

$$4e^{2x} = 16$$

$$e^{2x} = 4$$

$$lne^{2x} = ln4$$

$$2x = ln4$$

$$x = \frac{ln4}{2} \approx .693$$

5



$$e^{2+x} - 8 = 14$$

$$5e^{2+x} = 22$$

$$e^{2+x} = \frac{22}{5}$$

$$\ln e^{x+2} = \ln \frac{22}{5}$$

$$x + 2 = \ln \frac{22}{5}$$

$$x = \ln \frac{22}{5} - 2 \approx -0.518$$

Solving Exponential Equations

Solve:

 $5(4^{3x-7}) - 6 = 15$ $5(4^{3x-7}) = 21$

From this point you have a couple of choices.

$$\log_4 \frac{21}{5} = 3x - 7$$

$$3x = \log_4 \frac{21}{5} + 7$$
$$x = \frac{1}{3} \left(\log_4 \frac{21}{5} + 7 \right) \qquad x = \frac{1}{3} \left(\frac{\ln \frac{21}{5}}{\ln 4} + 7 \right)$$



$$4^{3x-7} = \frac{21}{5}$$

$$\ln 4^{3x-7} = \ln \frac{21}{5}$$

$$(3x - 7)\ln 4 = \ln \frac{21}{5}$$

$$3x - 7 = \frac{\ln \frac{21}{5}}{\ln 4} \qquad x = \frac{1}{3} \left(\frac{\ln \frac{21}{5}}{\ln 4} + 7 \right)$$

Quadratic Form of the Exponential

- Some exponential equations are in the form of a quadratic equation.
 - 1. Substitute u for the exponential term (u substitution).
 - 2. Solve the quadratic equation for u.
 - 3. Substitute the exponential for u.
 - 4. Solve for the variable.






Quadratic Form of the Exponential

$$\sim$$
 Solve: $e^{2x} - 8e^{x} + 7 = 0$

Let
$$u = e^{x}$$

Then $u^{2} = (e^{x})^{2} = e^{2x}$
 $e^{2x} - 8e^{x} + 7 = 0 \implies u^{2} - 8u + 7 = 0$
 $u^{2} - 8u + 7 = 0$
 $(u - 7)(u - 1) = 0$
 $u = 7 \text{ or } u = 1$





 \sim The solution set is {0, ln7}



Quadratic Form of the Exponential

$$\sim$$
 Solve: $2^{2x} + 2^{x} - 12 = 0$

Let
$$u = 2^{\times}$$

Then $u^2 = (2^{\times})^2 = 2^{2\times}$
 $2^{2\times} + 2^{\times} - 12 = 0 \implies u^2 + u - 12 = 0$
 $u^2 + u - 12 = 0$
 $(u - 3)(u + 4) = 0$
 $u = 3 \text{ or } u = -4$



 $ln2^{x} = ln3$ xln2 = ln3


$$x = \frac{\ln 3}{\ln 2} \approx 1.5850$$

The solution set is $\left\{\frac{\ln 3}{\ln 2}\right\}$





The graph $y = e^{2x} - 8e^{x} + 7$ and find the zeros.



Set your left \forall 2:Zero boundary by moving the cursor.



There is your problem. I will insist on exact solutions. Your calculator suggests a zero at about 1.94591.

 \sim Calculate $\ln(7)$

really and algebraically.

Set your right boundary by moving the cursor.





Check your Results

- Logarithmic expressions are defined only for logs of positive real numbers.
 - Always check proposed solutions of a logarithmic equation in the original equation.

Exclude from the solution set any proposed solution that produces the logarithm of a negative number or the logarithm of O (Extraneous solutions).





Using the Definition of a Logarithm to Solve Logarithmic Equations

- Sometimes you can convert a log equation into an exponential equation that is easier to solve.
 - 1. Express the equation in the form $log_b M = x$.
 - 2. Use the definition of a log to rewrite the equation in exponential form. $\log_b M = x \Rightarrow b^{\times} = M$
 - 3. Solve for the variable.
 - 4. Check proposed solutions in the original equation. Include in the solution set only values for which M > 0.





Example: Solving Logarifhmic Equations



$$log_{2}(x-4) = 3$$


$$2^{3} = (x-4)$$

$$x = 12$$

$$log_{2}(12-4) = 3$$

• The solution set is {12}.





- $4\ln 3x = 8$
 - ln3x = 2
 - $e^2 = 3x$

$$x = \frac{e}{3}$$

$$4\ln\left(3\left(\frac{e^2}{3}\right)\right) = 4\ln e^2 = 4(2) = 8$$

The solution set is $\left\{\frac{e^2}{3}\right\}$




Example: Solving Logarithmic Equations



- $\log x + \log(x-3) = 1$
 - logx(x-3) = 1
 - x(x-3) = 10
 - $x^2 3x 10 = 0$
 - (x-5)(x+2) = 0
 - x = 5 or x = -2





log10 = 1

 \sim The solution set is $\{5\}$.



Example: Solving Logarithmic Equations

Solve:

log(5x+1) = log(2x+3) + log2	log ₂ (x-3
log(5x+1) = log2(2x+3)	$\log_2\left(\frac{x}{x}\right)$
5x+1 = 4x+6	x(x)
x = 5	X -
log(26) = log(13) + log2	$\mathbf{x}(\mathbf{x})$
	x ² -
The solution set is $\{5\}$.	2





 $g_2(x-3) + log_2(x) - log_2(x+2) = 2$ $\frac{x(x-3)}{x+2} = 2$ (x-8)(x+1)=0 $\frac{(x-3)}{(x+2)} = 4$ x = 8 or x = -1(x-3) = 4(x+2)-3x = 4x + 8 $\log_{2} 5 + \log_{2} 8 - \log_{2} 10 = 2$ $x^2 - 7x - 8 = 0$



Using the One-to-One Property of Logarithms to Solve Logarithmic Equations

Writing the equation as equal logarithms.

- 1. Express the equation in the form $log_b M = log_b N$.
- 2. Use the one-to-one property to rewrite the equation without logs. If $log_b M = log_b N$, then M = N.
- 3. Solve for the variable
- 4. Check the proposed solutions in the original equation. Include in the solution set only value form which M>O and N>O.





Example: Solving a Logarithmic Equation



$$\log_2(x-1) - \log_2(x+3) = \log_2\left(\frac{1}{x}\right)$$

$$\log_{2}\left(\frac{x-1}{x+3}\right) = \log_{2}\left(\frac{1}{x}\right) \qquad \qquad x^{2}-x = x$$


$$\frac{x-1}{x+3} = \frac{1}{x}$$

x(x-1) = x+3

x = 3 or x - 1







$$x^{2}-2x-3=0$$

 $\log_{2}(2)-\log_{2}(6)=\log_{2}(\frac{1}{3})$
 $(x+1)(x-3)=0$





Example: Solving a Logarithmic Equation



ln(x-3) = ln(7x-23) - ln(x+1) $\ln(x-3) = \ln\frac{7x-23}{x+1}$ $x-3=\frac{7x-23}{3}$ x + 1(x-3)(x+1) = 7x-23 $x^2-2x-3 = 7x-23$ $x^2 - 9x + 20 = 0$ (x-5)(x-4) = 0 x=5 or x=4





ln(5-3) = ln(7(5)-23) - ln(5+1) $\ln 2 = \ln 12 - \ln 6$ ln(4-3) = ln(7(4)-23) - ln(4+1)ln1 = ln5 - ln5

The solution set is $\{4,5\}$.



Solving Exponential Equations



- 1. Rewrite the equation in the form $b^m = b^n$.
 - Note; the bases (b) are equal.
 - $4^{5x-2} = 8^{2x+3} \implies 2^{2(5x-2)} = 2^{3(2x+3)}$

2(5x-2) = 3(2x+3)

2. Take the ln (or log) of both sides of the equation.

 $6^{2x} = 1024 \implies \ln 6^{2x} = \ln 1024$

2xln6 = ln1024





Solving Logarithmic Equations



- 1. Rewrite the logarithmic equation in exponential form.
 - $\log_7(2x-5) = 11 \implies 7^{11} = 2x-5$
- 2. Simplify the logarithmic equation to $log_b m = log_b n$. lnx + ln(2x-1) = ln(5x-3) - ln2x

$$\mathbf{\sqrt{\mathbf{x}}} = \ln \frac{5x - 3}{2x}$$







Strategies for Solving Exponential and Logarithmic Equations

- **1.** Rewrite the original equation in a form that allows the use of the One-to-One Properties of exponential or logarithmic functions.
- **2.** Rewrite an *exponential* equation in logarithmic form and apply the Inverse Property of logarithmic functions.
- **3.** Rewrite a *logarithmic* equation in exponential form and apply the Inverse Property of exponential functions.



Example: Application

How long, to the nearest tenth of a year, will it take \$1000 to grow to \$3600 at 8% annual interest compounded quarterly?

 $A = P \left(1 + \frac{r}{n} \right)^{nr}$ $3600 = 1000 \left(1 + \frac{.08}{4}\right)^{47}$ $3.6 = (1 + .02)^{47}$ $\ln 3.6 = \ln (1.02)^{4t}$ $\ln 3.6 = 4t \ln(1.02)$

It will take about 16.2 years for \$1000 to become \$3600.



$$4t = \frac{\ln 3.6}{\ln 1.02}$$
$$t = \frac{\ln 3.6}{4\ln 1.02} \approx 16.17$$

